



DR. SAHAR DADKHAHFAR (Orcid ID : 0000-0003-4058-2562)

Article type : Letter to Editor

Skin Reactions to Non-glove Personal Protective Equipment: An Emerging Issue in the COVID-19 Pandemic

Mehdi Gheisari ^{1,+}, MD; Farnaz Araghi ^{1,+}, MD; Hamideh Moravvej ¹, MD;
Mohammadreza Tabary^{2,*}, MD; Sahar Dadkhahfar^{1,*}, MD

1. Skin Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran
2. School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

+ The first two authors had equal contributions

* Corresponding authors :

1. Sahar Dadkhahfar

e-mail: Sahar.dadkhahfar@gmail.com

Skin Research Center, Shahid Beheshti University of Medical Sciences, Shohada-e Tajrish Hospital, Shahr-dari St, 1989934148, Tehran, Iran.

Tel: +98-21-22741507 **Fax:** +98-21-22744393

2. Mohammadreza Tabary

e-mail: Mohammadrezatabary@gmail.com

School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/JDV.16492](https://doi.org/10.1111/JDV.16492)

This article is protected by copyright. All rights reserved

Tel: +98-21-22834154 **Fax:** +98-21-22834154

Running title: Personal Protective Equipment and Skin

Acknowledgment: The patients in this manuscript have given written informed consent to publication of their case details

Conflict of Interest: None to declare

Keywords: Personal protective equipment; skin reaction; COVID-19

Word Count: 608

Figure Count: 1

To Editor,

Protecting healthcare workers (HCWs) is crucial during Corona Virus Disease 2019 pandemic and requires wearing personal protective equipment (PPE) [1]. While most of the studies have focused on the skin reactions caused by gloves, other PPE such as gowns, respirator masks, face shields and goggles are also worn by HCWs for long hours during the current epidemic and skin irritations caused by these equipment may cause discouragement of health workers from using them [2]. In this letter we have focused on the reaction caused by non-glove PPE.

The N95 respirator masks are made of polypropylene fabric processed by a non-woven technology and should fit tightly to the face to be effective [3]. The study by Foo and colleagues revealed that 35.5% of the staff who used N95 masks regularly experienced acne, facial dermatitis and pigmentation of nasal bridge, cheeks and chin. In this study, acne was one of the most prevalent skin reactions related to the use of N95 respirator masks [3]. The dermatitis that often presented with pruritic skin lesions was mostly irritant type but allergic contact dermatitis (ACD) due to adhesives or other parts of the respirator masks such as rubber straps and metal clips was also reported [3]. Several factors including humidity, warm environment and occlusion due to local pressures could explain the exacerbation of these conditions [3]. In another study by Donovan *et al.*, on the possible N95 mask reactions during the SARS epidemic in Toronto, urticarial facial eruption was reported in 3 patients, dermatitis in 5 patients and acute respiratory symptoms without skin lesions in 2 patients [4]. Pressure effect on the nose have also been reported as one of the 15 delphi measures that discouraged HCWs to use N95 respirator masks [5].

Goggles, have been used routinely to protect HCWs against highly infectious diseases related to exposure to contaminated body fluids [6]. Heat and dehydration were major complications of both goggles and face shields application during the Ebola outbreak [7]. Other dermatologic side-effects such as pressure injury, contact dermatitis, urticaria, xerosis and aggravation of underlying dermatosis might occur due to the impairment of the skin integrity during mechanical trauma of goggles [8]. A study by Lan *et al.*, [2] revealed that 87.9% of HCWs, who were wearing goggles for more than 6 hours, developed skin reactions on their nasal bridge. Skin reactions such as acne, ACD, and irritant contact dermatitis (ICD) were mentioned following the use of goggles in HCWs. Occlusion and friction were mentioned as the underlying mechanism [6].

Wearing gowns and coveralls may cause heat stress and dehydration [7]. Skin reactions due to the clothing, which are made of natural and synthetic untreated fabrics, are rare [9]. However, additive chemicals and dye fibers might be the main reason of ICD and ACD [9]. Skin dermatoses are mostly developed where the gowns adhere tightly to the skin [10]. Friction, moisture and warmth of those regions might increase the risk of ACD[10]. In the study by *Foo et al.*, 4 (1.6%) out of 258 cases developed adverse skin reactions related to the repetitive wearing of disposable gowns for average time of 6.2 hours during a mean period of 8.8 months in the SARS epidemic in Singapore [3]. Itching and wrist rashes were the most frequent reactions, while pruritus without skin lesions was also observed in one case [3]. In Toronto SARS epidemic, there were reports of developing ACD due to the reaction to formaldehyde textiles and resin in gowns [10]. Avoiding over-tight gowns and sufficient ingestion of liquids are of paramount importance for HCWs to preserve a balance between self-protection and the ability to care for patients efficiently, while wearing PPE. Skin reactions to personal protective equipment and management strategies are depicted in Figure 1.

Figure legend:

Figure 1. Adverse skin reaction to personal protective equipment and management strategies

Reference:

- [1] Zhou P, Huang Z, Xiao Y, Huang X, Fan X-G. Protecting Chinese Healthcare Workers While Combating the 2019 Novel Coronavirus. *Infection Control & Hospital Epidemiology*. 2020; 1-4.
- [2] Lan J, Song Z, Miao X, Li H, Li Y, Dong L, et al. Skin damage and the risk of infection among healthcare workers managing coronavirus disease-2019. *Journal of the American Academy of Dermatology*. 2020.
- [3] Foo CCI, Goon ATJ, Leow YH, Goh CL. Adverse skin reactions to personal protective equipment against severe acute respiratory syndrome—a descriptive study in Singapore. *Contact dermatitis*. 2006;55; 291-294.
- [4] Donovan J, Kudla I, Holness LD, Skotnicki-Grant S, Nethercott JR. Skin reactions following use of N95 facial masks. *Dermatitis*. 2007;18; 104.
- [5] Honarbakhsh M, Jahangiri M, Farhadi P. Effective factors on not using the N95 respirators among health care workers: Application of Fuzzy Delphi and Fuzzy Analytic Hierarchy Process (FAHP). *Journal of Healthcare Risk Management*. 2017;37; 36-46.
- [6] Bhoirul B, Lecamwasam K, Wilkinson M, Latheef F, Stocks SJ, Agius R, et al. A review of non-glove personal protective equipment-related occupational dermatoses reported to EPIDERM between 1993 and 2013. *Contact Dermatitis*. 2019;80; 217-221.
- [7] Den Boon S, Vallenias C, Ferri M, Norris SL. Incorporating health workers' perspectives into a WHO guideline on personal protective equipment developed during an Ebola virus disease outbreak. *F1000Research*. 2018;7.
- [8] Yan Y, Chen H, Chen L, Cheng B, Diao P, Dong L, et al. Consensus of Chinese experts on protection of skin and mucous membrane barrier for healthcare workers fighting against coronavirus disease 2019. *Dermatologic Therapy*. 2020; e13310.
- [9] Fowler JF. Formaldehyde as a textile allergen. *Curr Probl Dermatol*. 2003;31; 156-165.
- [10] Donovan J, Skotnicki-Grant S. Allergic contact dermatitis from formaldehyde textile resins in surgical uniforms and nonwoven textile masks. *Dermatitis*. 2007;18; 40-44.



jd_v_16492_f1.jpg